



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Cereal powdery mildew effectors: a complex toolbox for an obligate pathogen

Bourras, Salim ; Praz, Coraline R ; Spanu, Pietro D ; Keller, Beat

Abstract: Highlights • Cereal powdery mildews have evolved a large repertoire of candidate effector proteins. • These effectors are encoded within size variable and highly diversified gene families. • Mildew effectors can act as virulence factors, Avr genes and Avr suppressors. • Their mode of action provides a molecular basis for the Zig-Zag evolutionary model.

DOI: <https://doi.org/10.1016/j.mib.2018.01.018>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-160637>

Journal Article

Accepted Version

Originally published at:

Bourras, Salim; Praz, Coraline R; Spanu, Pietro D; Keller, Beat (2018). Cereal powdery mildew effectors: a complex toolbox for an obligate pathogen. *Current Opinion in Microbiology*, 46:26-33.

DOI: <https://doi.org/10.1016/j.mib.2018.01.018>

Dear author,

Please note that changes made in the online proofing system will be added to the article before publication but are not reflected in this PDF.

We also ask that this file not be used for submitting corrections.



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Microbiology

Cereal powdery mildew effectors: a complex toolbox for an obligate pathogen

Salim Bourras¹, Coraline R Praz¹, Pietro D Spanu² and Beat Keller¹

Cereal powdery mildews are major pathogens of cultivated monocot crops, and all are obligate biotrophic fungi that can only grow and reproduce on living hosts. This lifestyle is combined with extreme host specialization where every mildew subspecies (referred to as *forma specialis*) can only infect one plant species. Recently there has been much progress in our understanding of the possible roles effectors play in this complex host–pathogen interaction. Here, we review current knowledge on the origin, evolution, and mode of action of cereal mildew effectors, with a particular focus on recent advances in the identification of *bona fide* effectors and avirulence effector proteins from wheat and barley powdery mildews.

Addresses

¹ Department of Plant and Microbial Biology, University of Zürich, Zollikerstrasse 107, 8008 Zürich, Switzerland

² Department of Life Sciences, Imperial College London, Imperial College Road, SW7 2AZ London, United Kingdom

Corresponding author: Bourras, Salim (s.bourras@botinst.uzh.ch)

Current Opinion in Microbiology 2018, 46:xx–yy

This review comes from a themed issue on **Host–microbe interactions: fungi**

Edited by **Ralph Dean** and **Thierry Rouxel**

<http://dx.doi.org/10.1016/j.mib.2018.01.018>

1369–5274/© 2018 Published by Elsevier Ltd.

Introduction

Powdery mildews are agronomically important fungal pathogens infecting a wide range of monocot and dicot crops. Cereal powdery mildew diseases are caused by only one species, *Blumeria graminis*, which can be divided into several subspecies, corresponding to highly specialized pathogens infecting only one specific crop species. These are sometimes referred to as '*forma specialis*' (f.sp., literally forms belonging to one host species). For instance, *Blumeria graminis tritici* (*B.g. f. sp. tritici*), *B.g. hordei*, and *B.g. secalis*, can only grow on wheat, barley, or rye respectively [1[•],2,3]. All cereal mildews are obligate biotrophs, meaning that they can only grow and reproduce on living host

tissue. Considering the fact that this lifestyle is combined with extreme host specialization, it was proposed that highly complex and yet poorly understood mechanisms regulate host–pathogen interactions in cereal mildews [4^{••},5^{••}]. Whole genome sequencing of the wheat and barley powdery mildews revealed on the one hand a drastic reduction of the gene content compared to other ascomycete pathogens, and on the other hand an expansion of the putative effector gene complement [4^{••},5^{••}]. In this context, we propose that cereal powdery mildews provide a highly informative system to study the molecular role of effectors in pathogen virulence and host adaptation, based on a case of extreme host specialization. This review aims at providing a concise overview of current knowledge in effector biology of cereal mildews. In particular, we summarise advances in effector gene identification and functional characterization. Finally, possible ways for accelerating effector gene function discovery in cereal mildews will be discussed.

Genome-wide identification of candidate effectors in cereal mildew genomes

In powdery mildews, putative effectors have been identified using whole genome sequencing, and large-scale analysis of the fungal proteome, based on *a priori* criteria defining groups of protein coding genes that eventually constituted the putative effectors. Thus, Candidate Secreted Effector Proteins (CSEPs) from wheat and barley powdery mildew were defined as predicted secreted proteins (i.e. including an N-terminal 'signal peptide') that did not have trans-membrane domains, and did not have homologues in non-mildew fungi [4^{••},5^{••}]. Based on these criteria, initial sets of 472 and 437 CSEPs were identified in the barley and wheat powdery mildew genomes, respectively. There have been continuous efforts to improve effector gene prediction in mildews which has resulted in the identification of larger sets of 722 and 734 CSEPs in the barley and wheat powdery mildew genomes, respectively [6^{••},7].

In addition to genome information, powdery mildew effectors were also mined in the proteome of *B.g. hordei*, by identifying proteins that are found in fractions enriched with isolated haustoria [8] or in infected barley epidermis devoid of epiphytic fungal material [9,10]. In the latter work, to generate a group of proteins named *Blumeria Effector Candidate* (BEC) proteins, the criterion of excluding homologs present in non-mildew fungi

2 Host

was not used. This has led to the identification of effector proteins such as BEC1005 and BEC1019, which are necessary for full virulence in barley powdery mildew, and are broadly conserved in ascomycete fungi [11,12]. Interestingly, there is nearly complete overlap between the set of barley powdery mildew CSEPs and the BECs; that is, there are only five BECs which are not included in the CSEP set [13].

Origin and evolution of mildew effectors

The Erysiphales, that is the fungi causing powdery mildews, are an ancient monophyletic group which are estimated to have originated within the Leotiomyces over 120 million years ago (Mya) [14]. The closest sister group is the non-pathogenic Myxotrichaceae, while all extant Erysiphales are obligate biotrophic pathogens of plants. It is reasonable to assume that effector proteins played essential roles in the early evolution of the powdery mildew fungi. One good example are the genes encoding for effector proteins with a predicted fungal ribonuclease-like (RNase-like) three dimensional structure similar to the ribonuclease T1 from *Aspergillus phoenicis* [6^{••}]. The genes encoding the so-called RNase-like effectors have a single intron in a strictly conserved position [15]. Remarkably, both RNase-like effectors and the conserved intron are present in *Erysiphe necator* the agent of grapevine powdery mildew, which is thought to have diverged from the cereal mildew fungi in the late Cretaceous, over 70 Mya [Spanu and Dry, unpublished results]. Therefore, it is likely that this class of mildew effectors accounting for ca. 10–15% of all predicted mildew CSEPs and represented in six out of the 20 largest mildew effector families [6^{••},7] derived from a single ancestor similar to the canonical fungal RNase T1 [16].


In a recent study, Menardo and colleagues found a new class of CSEPs with structural homologies with the MD2-related lipid-recognition (ML) domain [7], which is predicted to be involved in binding to specific lipids (IPR003172) [17]. Notably, the gene family encoding for ML-like CSEPs is conserved in distinctly different lineages of grass powdery mildews, hinting this novel class of mildew effectors may also be derived from a common ancestor [7].

Cereal powdery mildew effector families are under stronger diversifying selection than non-effector genes, and show extreme levels of sequence variation and gene turnover [7], while the predicted structures appear conserved (Figures 1 and 2a) [18]. Essentially, only the protein sequence encoding the predicted signal peptide is highly conserved among members of the same family. Immediately after the putative cleavage site, sequence homology is reduced to a conserved variant of the YxC motif [6^{••}], and the position of a few amino acids including residues of putative structural importance such as cysteines and prolines (Figure 2a). This high level of sequence divergence might be a result of a strong diversifying selection pressure imposed by the host immune system. Indeed, several RNase-like effectors from wheat, barley, and rye powdery mildews interact with nucleotide-binding leucine-rich repeat (NLR) immune receptors from the host, or suppress such a recognition, suggesting they are likely under selection to evolve [19^{••},20,21^{••},22^{••}].

Mode of action of cereal mildew effectors


The first insights into the mode of action of cereal mildew effectors were largely based on transcriptomics and

Figure 1



Predicted three-dimensional models of powdery mildew RNase-like effectors. Predicted three-dimensional protein folds are based on the crystal structure of the *Aspergillus phoenicis* Ribonuclease T1 (a). The wheat powdery mildew CSEP AVRPM2 (b), and the barley powdery mildew CSEP AVR_{A13} (c) are depicted. AVRPM2 and AVR_{A13} are encoded within the same effector gene family, and they are recognized by the evolutionary unrelated wheat PM2 and the barley MLA13 immune receptors, respectively.

Figure 2



Common features of cereal powdery mildew effector gene families. **(a)** Protein sequence alignment of the AVRPM2 effector family members from *B.g. tritici* (red bars) and *B.g. hordei* (black bars). Protein sequences corresponding to the mildew AVR proteins AVRPM2 and AVR_{A13}, and the HIGS validated mildew *bona fide* effectors BEC1054 and BEC1011, are highlighted with prominent horizontal bars. The N-terminal predicted signal peptide, the conserved YxxC motif, as well as the most conserved residues are framed and annotated accordingly. Region under diversifying selection pressure is highlighted. **(b)** RNA-sequencing based assessment of gene expression levels across the *B.g. tritici* members of the AvrPm3^{a2/f2} (left panel), AvrPm2 (middle panel) effector gene families in the wheat powdery mildew isolate Bgt-96224, and the SvrPm3^{a1/f1} effector gene family in the wheat mildew isolate Bgt-94202. Expression levels are given in reads per kilobase per million mapped reads (RPKM). Every bar corresponds to the expression of one family member at the wheat mildew haustorial stage (48 hour post infection), during compatible interaction between *B.g. tritici* and the susceptible wheat cultivar 'Chinese Spring' (Coraline Praz, unpublished data). Effectors in the first quartile are transcriptionally inactive and constitute a proposed group of latent effectors. By contrast, effectors in the fourth quartile are always highly expressed and encode for active Avr and Svr genes whose position in the plots is indicated with an arrow.

proteomics approaches in *B.g. hordei* [8–10,13]. In particular, RNA-seq monitoring of barley powdery mildew transcripts during the early stages of host infection revealed a two-step mode of action: a first wave of CSEP

transcripts accumulated during host cell entry (12 hour), and a second wave of transcripts accumulated at the stage of haustorium formation (24 hour) [23[•]]. Similarly, high induction of CSEPs at the haustorial stage (48 hour) is

4 Host

also observed in wheat powdery mildew (Figure 2b; Coraline Praz, unpublished data). These results substantiate the importance of candidate effectors for mildew virulence, and suggest there are different subsets of CSEPs fulfilling distinct biological functions depending on the developmental stages of the fungus during host colonization. Indeed, Host Induced Gene Silencing (HIGS) [24^{••}] of 21 individual barley powdery mildew effector genes resulted in significant reduction of host penetration and haustorium formation, further supporting the essential contribution of mildew CSEPs to the establishment of host infection [11,25[•],26[•],27]. Pliego and colleagues also showed that HIGS downregulation of 50 haustorially expressed barley mildew effectors resulted in highly variable effects on fungal virulence, ranging from a significant increase to a significant reduction of haustorium formation depending on the targeted effector gene [11]. This data suggests that some CSEPs are probably dispensable, while others such as BEC1054 and BEC1011, whose HIGS downregulation resulted in 60–70% reduction of haustorium formation, are acting as bona fide (i.e. true) effectors that are essential for mildew virulence [11].

There is evidence that mildew effectors interfere with components of host basal metabolism and host immunity, with prominent examples being CSEP0055 [25[•]], BEC3, BEC4 [28[•]], CSEP0105, CSEP0162 [26[•]], and BEC1054 [29[•]] from *B.g. hordei*, and *SvrPm3^{al/f1}* [19^{••},20] from *B.g. tritici* (Figure 3). Ahmed and colleagues showed that the sequence unrelated CSEP0105 and CSEP0162 both interact with the stress related small heat shock protein chaperones HSP16.9 and HSP17.5 from barley [26[•]]. We propose a possible mode of action of mildew CSEPs based on functional redundancy among effectors (Figure 3). In an approach combining pull-down assays from barley protein extracts and experimental validation by yeast-2-hybrid, Pennington and colleagues showed that BEC1054 physically interacts with a barley pathogen-related-5 (PR5) protein plus three sequence unrelated proteins fulfilling distinctly different biochemical functions, including a glutathione-S-transferase (GST), a malate dehydrogenase (MDH), and an elongation factor 1 gamma protein (eEF1G) [29[•]]. It is unclear if these interactions are based on the presence a conserved protein motif in all four targets serving as a binding site for BEC1054, or if the effector protein itself carries several protein–protein interaction domains allowing specific binding to multiple targets. It is also unclear how interactions between BEC1054 and its barley targets are affecting the host's metabolism or immunity. Nevertheless, the experimental data from Pennington and colleagues suggests another possible mode of action of mildew CSEPs based on specific binding to several host targets. Here, the action of a single effector can possibly disturb several biochemically distinct proteins involved in distinctly different pathways, which we could refer to as 'Effector disturbance' (Figure 3). Finally, in wheat powdery mildew,


Bourras and colleagues provided genetic evidence that the SVRPM3^{A1/F1} suppressor of AVR (SVR) recognition, is a suppressor of the wheat PM3A and PM3F-mediated race-specific resistance of wheat, which has been functionally demonstrated in *Agrobacterium*-mediated transient expression assays in *Nicotiana benthamiana* [19^{••}]. There are only a few examples of SVR effectors from plant pathogens, with two other prominent examples being AVR1 from the tomato pathogen *Fusarium oxysporum* [30] and AVR1M4-7 from the oilseed rape pathogen *Leptosphaeria maculans* [31]. The mode of action of the SVRPM3^{A1/F1} RNase-like CSEP is unknown, and we propose this can be based either on a ribonuclease pseudoenzyme activity [32] or on an effector disturbance function targeting components of the PM3 resistance signaling.

There are 167 family clusters that have been identified across *B.g. tritici*, *B.g. hordei*, *B.g. on Lolium*, *B.g. avenae*, and *B.g. poae* [7], which may possibly encode a large diversity of biochemical functions with different modes of actions. There are also significant differences at the gene expression level within families, with some members showing very high levels of expression at the haustorial stage, while others seem to be transcriptionally extinct (Figure 2b; Coraline Praz, unpublished data). One possible scenario is that 'active' effectors are exposed to selective pressure from the host, and are at risk of being recognized by the plant immune system. In such cases, these effectors are transcriptionally suppressed and become 'latent'. If this were the case, they could represent a reservoir of effectors which can be reactivated to compensate for the loss of function, mutation, deletion, or downregulation of other active effectors imposed by selective pressure to escape host recognition. We speculate that this transcriptional plasticity allows the preservation of high levels of genetic diversity in effector families to provide cereal mildews with a potential for cultivar and host adaptation. We also speculate that the very high repeat content of the mildew genomes might contribute to such plasticity by providing additional layers of regulation at the epigenetic level [33,34].

Mildew effectors recognized by the plant immune system

Genes encoding CSEPs in plant pathogenic fungi may act as avirulence genes (*Avr*) recognized by the plant resistance (*R*) gene based immunity. In the recent years, the identification of CSEPs encoding AVR proteins in mildews has been greatly accelerated by increasing integration of next-generation sequencing and high-throughput genotyping technologies. Mildew *Avrs* have been thus identified using map-based cloning [19^{••},22^{••}], bulk-segregant-analysis (BSA) [19^{••},35], genome wide association studies (GWAS) [22^{••}], and RNAseq based GWAS [21^{••}]. This has allowed the cloning of four *Avr* effectors: the *B.g. tritici* *AvrPm3^{a2/f2}*, and *AvrPm2* which are recognized by

Figure 3



Contribution of cereal powdery mildew CSEPs to host-pathogen interactions. Three layers of plant-pathogen interactions commonly defined in the Zig-Zag evolutionary model are here represented. In this model, a subset of powdery mildew effector proteins (CSEP0105, CSEP0162, BEC1054, BEC3, and BEC4) are employed at suppressing the first layer of plant defenses commonly provided by PAMP-triggered immunity (PTI), or at manipulating the host cell to derive nutrients, thus resulting in a first layer of Effector-triggered susceptibility (ETS_1, upper panel). Here we highlight two possible modes of actions, namely 'Effector redundancy' and 'Effector disturbance' as well as functionally validated examples of effector host targets (see text for details). Another subset of effectors (AVR_{A1} , $AVRPM2$, AVR_{A13} , and $AVRPM3^{A2/F2}$), are probably also actively involved in virulence, but they are recognized by NLR receptors from the plant immune system (MLA1, PM2, MLA13, and PM3A/F, respectively), thus leading to Effector-triggered immunity (ETI, middle panel). These are basically canonical AVR-NLR interactions controlling race specific resistance in the host. Finally, a third subset of effectors ($SVRPM3^{A1/F1}$) are SVRs capable of suppressing such AVR-NLR mediated resistance thus providing a second layer of Effector-triggered susceptibility (ETS_2) based on the AVR-R-SVR model.

the wheat *Pm3alf*, and *Pm2* R genes, respectively [19^{••}, 22^{••}], and the *B.g. hordei* *Avr_{a1}*, *Avr_{a13}* which are recognized by the barley *Mla1* and *Mla13* R genes, respectively [21^{••}]; and one suppressor of *Avr* recognition, *SvrPm3^{a1/f1}* [19^{••}, 20]. *AvrPm2* and *Avr_{a13}* belong to the same effector gene family, and together with *Avr_{a1}* and *SvrPm3^{a1/f1}* they encode for RNase-like effectors

(Figures 1 and 2a). One exception so far is *AvrPm3^{a2/f2}*, the cognate *Avr* of the *Pm3a* and *Pm3f* resistance gene alleles from wheat, which does not encode for a ribonuclease-like protein [19^{••}]. Mildew *Avrs* are also members of size variable effector families, and all are among the most highly induced members upon infection (Figure 2b). This data suggests that members of the same

6 Host

effector gene family do not equally contribute to virulence, and that the most highly expressed ones are likely to be recognized by the plant immune system as avirulence factors.

Altogether, these results show that mildew effectors can act on the multiple layers of plant immunity described in the Zig-Zag model of plant–pathogen interactions [36,37] (Figure 3). Some effectors proteins such as BEC1054, may act as *bona fide* effectors suppressing PAMP-triggered immunity (PTI) and allowing successful host colonization through a first layer of effector-triggered susceptibility (ETS₁, Figure 3). The biochemical basis of such mode of action remains unclear since little is known about effector function in mildews. Other effectors such as AVRPM3^{A2/F2}, are canonical targets of plant immune receptors, whose recognition results in effector-triggered immunity (ETI, Figure 3). A recent study by McNally and colleagues revealed that *AvrPm3*^{a2/f2} is highly conserved in worldwide wheat mildew populations, suggesting this CSEP may possibly act as an important virulence factor on susceptible hosts [38]. Finally, another type of CSEPs such as SVRPM3^{A1/F1} are *bona fide* effectors that may act as SVRs suppressing ETI, allowing the pathogen to escape *R* gene recognition, and therefore achieving a second layer of ETS (ETS₂, Figure 3).

Future work on cereal mildew effectors

There is no transformation protocol for cereal powdery mildew fungi, nor is there any method for cultivation of these fungi on artificial media. These constraints impose the use of alternative approaches for the characterization of powdery mildew effectors based on heterologous plant, fungal, and bacterial expression systems. So far, studies combining HIGS, and effector targets screens (e.g. yeast-2-hybrid and pull-down assays) have succeeded in the functional validation of the effector function of several CSEPs, and in subsequent identification of several of their host targets. Other important and parallel efforts have resulted in the identification of CSEPs encoding mildew AVR; several AVR–NLR pairs have been functionally validated using transient expression assays in barley protoplasts, and *Agrobacterium*-mediated transient expression in barley and in the heterologous *N. benthamiana* host, three assays amenable to high-throughput screens.

On the host side, more than 60 powdery mildew resistance genes have been described in the wheat gene pool, many of them with different alleles [39–41]. Similarly, in barley there are many resistance genes known, the large *Mla* allelic series being the most prominent [42,43]. Therefore, we propose that the identification of the cognate CSEPs for many of these resistance proteins, combined with functional characterization of their effector function and virulence targets, will provide a unique biological opportunity to describe the network of

interactions between cereal powdery mildews and their hosts. It should also be possible to identify AVR encoding CSEPs based on common features of *Avr* genes in mildews (e.g. they are sequence polymorphic and highly expressed members of their effector gene family), which can be mined in the large number of available mildew genomes and transcriptomes. We also propose that the resolution of the three-dimensional structure of representative members of large CSEP families, can be a powerful tool to dissect effector protein functions based on structure-informed mutagenesis screens. Finally, the field of *Avr* population genetics is largely underexplored in cereal mildews, and we suggest the study of *Avr* diversity at the population level will be another powerful tool for understanding the evolutionary forces driving host–pathogen co-evolution in this agronomically important pathosystem.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Menardo F, Praz CR, Wyder S, Ben-David R, Bourras S, Matsumae H, McNally KE, Parlange F, Riba A, Roffler S *et al.*: **Hybridization of powdery mildew strains gives rise to pathogens on novel agricultural crop species.** *Nat Genet* 2016, **48**:201–205. This study shows that triticale powdery mildew, a new pathogen that has recently emerged on the triticale crop, arose from a hybridization event between wheat and rye powdery mildews. The authors show that the triticale mildew genome is a mosaic of segments from the wheat and rye donors. These conclusions are based on the sequencing and analysis of 46 powdery mildew isolates covering several *formae speciales*.
2. Menardo F, Wicker T, Keller B: **Reconstructing the evolutionary history of powdery mildew lineages (*Blumeria graminis*) at different evolutionary time scales with NGS data.** *Genome Biol Evol* 2017, **9**:446–456.
3. Troch V, Audenaert K, Wyand RA, Haesaert G, Höfte M, Brown JKM: **Formae speciales of cereal powdery mildew: close or distant relatives?** *Mol Plant Pathol* 2014, **15**:304–314.
4. Spanu PD, Abbott JC, Amselem J, Burgis TA, Soanes DM, Stuber K, Loren van Themaat EV, Brown JKM, Butcher SA, Gurr SJ *et al.*: **Genome expansion and gene loss in powdery mildew fungi reveal tradeoffs in extreme parasitism.** *Science* 2010, **330**:1543–1546. This paper describes the barley powdery mildew draft reference genome, as well as the analysis and comparison with two powdery mildews pathogens of dicotyledonous plants. Evidence is provided for a massive retrotransposon proliferation in the mildew genomes, coupled with a drastic loss of genes encoding enzymes of primary and secondary metabolism, carbohydrate-active enzymes, and transporters.
5. Wicker T, Oberhaensli S, Parlange F, Buchmann JP, Shatalina M, Roffler S, Ben-David R, Doleel J, Imková H, Schulze-Lefert P *et al.*: **The wheat powdery mildew genome shows the unique evolution of an obligate biotroph.** *Nat Genet* 2013, **45**:1092–1096. This paper describes the wheat powdery mildew draft reference genome, the resequencing of three additional isolates, and comparative analyses with the barley powdery mildew genome. This study provides evidence that the mildew genomes are mosaics of ancient haplogroups that existed before wheat domestication.
6. Pedersen C, van Themaat EVL, McGuffin LJ, Abbott JC, Burgis TA, Barton G, Bindschedler LV, Lu X, Maekawa T, Wessling R *et al.*: **Structure and evolution of barley powdery mildew effector candidates.** *BMC Genomics* 2012, **13**:694.

- In this work, genomic, transcriptomic and proteomic analyses as well as structural prediction methods were combined to identify and characterize the members of the CSEPs superfamily. This pioneering work led to the identification of RNase-like CSEPs, the largest class of effector proteins in mildew.
7. Menardo F, Praz CR, Wicker T, Keller B: **Rapid turnover of effectors in grass powdery mildew (*Blumeria graminis*). *BMC Evol Biol* 2017. in press.**
 8. Godfrey D, Zhang Z, Saalbach G, Thordal-Christensen H: **A proteomics study of barley powdery mildew haustoria. *Proteomics* 2009, 9:3222–3232.**
 9. Bindschedler LV, Burgis TA, Mills DJS, Ho JTC, Cramer R, Spanu PD: **In planta proteomics and proteogenomics of the biotrophic barley fungal pathogen *Blumeria graminis* f. sp. *hordei*. *Mol Cell Proteomics* 2009, 8:2368–2381.**
 10. Bindschedler LV, McGuffin LJ, Burgis TA, Spanu PD, Cramer R: **Proteogenomics and in silico structural and functional annotation of the barley powdery mildew *Blumeria graminis* f. sp. *hordei*. *Methods* 2011, 54:432–441.**
 11. Pliego C, Nowara D, Bonciani G, Gheorghe DM, Xu R, Surana P, Whigham E, Nettleton D, Bogdanove AJ, Wise RP et al.: **Host-induced gene silencing in barley powdery mildew reveals a class of ribonuclease-like effectors. *Mol Plant–Microbe Interact* 2013, 26:633–642.**
 12. Whigham E, Qi S, Mistry D, Surana P, Xu R, Fuerst G, Pliego C, Bindschedler LV, Spanu PD, Dickerson JA et al.: **Broadly conserved fungal effector BEC1019 suppresses host cell death and enhances pathogen virulence in powdery mildew of barley (*Hordeum vulgare* L.). *Mol Plant–Microbe Interact* 2015, 28:968–983.**
 13. Bindschedler LV, Panstruga R, Spanu PD: **Mildew-omics: how global analyses aid the understanding of life and evolution of powdery mildews. *Front Plant Sci* 2016, 7.**
 14. Takamatsu S: **Origin and evolution of the powdery mildews (ascomycota, erysiphales). *Mycoscience* 2013, 54:75–86.**
 15. Spanu PD: **Cereal immunity against powdery mildews targets RNase-Like Proteins associated with Haustoria (RALPH) effectors evolved from a common ancestral gene. *New Phytol* 2017, 213:969–971.**
 16. Yoshida H: **The ribonuclease T1 family. *Methods Enzymol* 2001, 341:28–41.**
 17. Inohara N, Nuez G: **ML – a conserved domain involved in innate immunity and lipid metabolism. *Trends Biochem Sci* 2002, 27:219–221.**
 18. Bourras S, McNally KE, Müller MC, Wicker T, Keller B: **Avirulence genes in cereal powdery mildews: the gene-for-gene hypothesis 2.0. *Front Plant Sci* 2016, 7:241.**
 19. Bourras S, McNally KE, Ben-David R, Parlange F, Roffler S, Praz CR, Oberhaensli S, Menardo F, Stirnweis D, Frenkel Z et al.: **Multiple avirulence loci and allele-specific effector recognition control the Pm3 race-specific resistance of wheat to powdery mildew. *Plant Cell* 2015, 27:2991–3012.**
This study reports the cloning of the first mildew effector gene encoding for a typical AVR protein, *AvrPm3^{2/12}*; and the identification of *SvrPm3^{3/11}*, the first suppressors of AVR recognition in mildews. The authors propose an extension of the Flor's gene-for-gene model based on a three component genetic interaction between avirulence and suppressor genes from the pathogen, and a resistance gene from the host.
 20. Parlange F, Roffler S, Menardo F, Ben-David R, Bourras S, McNally KE, Oberhaensli S, Stirnweis D, Buchmann G, Wicker T et al.: **Genetic and molecular characterization of a locus involved in avirulence of *Blumeria graminis* f. sp. *tritici* on wheat Pm3 resistance alleles. *Fungal Genet Biol* 2015, 82:181–192.**
 21. Lu X, Kracher B, Saur IML, Bauer S, Ellwood SR, Wise R, Yaeno T, Maekawa T, Schulze-Lefert P: **Allelic barley MLA immune receptors recognize sequence-unrelated avirulence effectors of the powdery mildew pathogen. *Proc Natl Acad Sci* 2016, 113: E6486–E6495.**
This study reports the cloning of two typical avirulence effector genes from barley powdery mildew, *Avr_{a1}* and *Avr_{a13}*. These Avr's were identified with a novel approach using RNA sequencing data as basis for GWAS in only 17 isolates of barley mildew. The authors show that *Avr_{a1}* and *Avr_{a13}* are sequence and phylogenetically unrelated, yet they are recognized by two alleles of the same barley NLR immune receptor.
 22. Praz CR, Bourras S, Zeng F, Sánchez-Martín J, Menardo F, Xue M, Yang L, Roffler S, Böni R, Herren G et al.: **AvrPm2 encodes an RNase-like avirulence effector which is conserved in the two different specialized forms of wheat and rye powdery mildew fungus. *New Phytol* 2017, 213:1301–1314.**
This study reports the cloning of *AvrPm2* from wheat powdery mildew, the cognate Avr of the *Pm2* resistance gene. The authors show that this Avr belongs to an effector family containing functionally validated effectors and an additional Avr gene from barley powdery mildew, *Avr_{a13}*, and all encode for RNase-like proteins. This study substantiates the important role this class of effector proteins is playing in host–mildew interactions.
 23. Hacquard S, Kracher B, Maekawa T, Vernaldi S, Schulze-Lefert P, Ver Loren van Themaat E: **Mosaic genome structure of the barley powdery mildew pathogen and conservation of transcriptional programs in divergent hosts. *Proc Natl Acad Sci* 2013, 110:E2219–E2228.**
This is the first detailed transcriptome analysis of early powdery mildew pathogenesis on barley and immunocompromised Arabidopsis. This work describes a two steps (waves) mode of action of mildew CSEPs, and reveals strong conservation of gene regulation in barley powdery mildew on both hosts despite ~200 million years of reproductive isolation.
 24. Nowara D, Gay A, Lacomme C, Shaw J, Ridout C, Douchkov D, Hensel G, Kümlehn J, Schweizer P: **HIGS: host-induced gene silencing in the obligate biotrophic fungal pathogen *Blumeria graminis*. *Plant Cell* 2010, 22:3130–3141.**
This work describes one of the most important tools available for functional genomics in powdery mildews. This technique has allowed the functional validation of the effector function of several mildew CSEPs.
 25. Zhang WJ, Pedersen C, Kwaaitaal M, Gregersen PL, Mørch SM, Hanisch S, Kristensen A, Fuglsang AT, Collinge DB, Thordal-Christensen H: **Interaction of barley powdery mildew effector candidate CSEP0055 with the defence protein PR17c. *Mol Plant Pathol* 2012, 13:1110–1119.**
This study provides evidence that powdery mildew CSEPs can target defence proteins from the host.
 26. Ahmed AA, Pedersen C, Schultz-Larsen T, Kwaaitaal M, Jørgensen HJL, Thordal-Christensen H: **The barley powdery mildew candidate secreted effector protein CSEP0105 inhibits the chaperone activity of a small heat shock protein. *Plant Physiol* 2015, 168:321–333.**
This study provides evidence that sequence unrelated powdery mildew CSEPs can target the same host protein, thus providing evidence for a possible mode of action of mildew effectors based on functional redundancy.
 27. Ahmed AA, Pedersen C, Thordal-Christensen H: **The barley powdery mildew effector candidates CSEP0081 and CSEP0254 promote fungal infection success. *PLoS One* 2016, 11.**
 28. Schmidt SM, Kuhn H, Micali C, Liller C, Kwaaitaal M, Panstruga R: **Interaction of a *Blumeria graminis* f. sp. *hordei* effector candidate with a barley ARF-GAP suggests that host vesicle trafficking is a fungal pathogenicity target. *Mol Plant Pathol* 2014, 15:535–549.**
This study provides evidence that host proteins involved in basal functions are targeted by mildew CSEPs, suggesting mildew effectors can possibly reprogram the host cellular machinery.
 29. Pennington HG, Gheorghe DM, Damerum A, Pliego C, Spanu PD, Cramer R, Bindschedler LV: **Interactions between the powdery mildew effector BEC1054 and barley proteins identify candidate host targets. *J Proteome Res* 2016, 15:826–839.**
This study provides evidence that mildew CSEPs can target several sequence unrelated and functionally distinct proteins in the host, thus providing evidence for a possible mode of action of mildew effectors based on effector disturbance.
 30. Houterman PM, Cornelissen BJC, Rep M: **Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* 2008, 4:e1000061.**
 31. Plissonneau C, Daverdin G, Ollivier B, Blaise F, Degraeve A, Fudal I, Rouxel T, Balesdent MH: **A game of hide and seek between avirulence genes *AvrLm4-7* and *AvrLm3* in *Leptosphaeria maculans*. *New Phytol* 2016, 209:1613–1624.**

8 Host

- 504 32. Adrain C, Freeman M: **New lives for old: evolution of**
505 **pseudoenzyme function illustrated by iRhoms.** *Nat Rev Mol Cell*
506 *Biol* 2012, **13**:489–498.
- 507 33. Soyer JL, El Ghalid M, Glaser N, Ollivier B, Linglin J, Grandaubert J,
508 Balesdent M-H, Connolly LR, Freitag M, Rouxel T *et al.*:
509 **Epigenetic control of effector gene expression in the plant**
510 **pathogenic fungus *Leptosphaeria maculans*.** *PLoS Genet* 2014,
10:e1004227.
- 511 34. Gijzen M, Ishmael C, Shrestha SD: **Epigenetic control of**
512 **effectors in plant pathogens.** *Front Plant Sci* 2014, **5**.
- 513 35. Takagi H, Abe A, Yoshida K, Kosugi S, Natsume S, Mitsuoka C,
514 Uemura A, Utsushi H, Tamiru M, Takuno S *et al.*: **QTL-seq: rapid**
515 **mapping of quantitative trait loci in rice by whole genome**
516 **resequencing of DNA from two bulked populations.** *Plant J*
2013, **74**:174–183.
- 517 36. Jones JDG, Dangl JL: **The plant immune system.** *Nature* 2006,
518 **444**:323–329.
- 519 37. Hein I, Gilroy EM, Armstrong MR, Birch PRJ: **The zig-zag-zig**
520 **in oomycete–plant interactions.** *Mol Plant Pathol* 2009,
521 **10**:547–562.
- 522 38. McNally KE, Menardo F, Lüthi L, Praz CR, Müller MC, Kunz L, Ben-
David R, Chandrasekhar K, Dinor A, Cowger C *et al.*: **Distinct**
523 **domains of the AVRPM3A2/F2 avirulence protein from wheat**
powdery mildew are involved in immune receptor recognition
and putative effector function. *New Phytol* 2018 <http://dx.doi.org/10.1111/nph.15026>.
- 524
525
526
39. McIntosh R, Yamazaki Y: **Catalogue of gene symbols for wheat.**
Int Wheat Genet Symp 2008, **4**:1–197.
- 527
528
40. Bhullar NK, Zhang Z, Wicker T, Keller B: **Wheat gene bank**
accessions as a source of new alleles of the powdery mildew
resistance gene Pm3: a large scale allele mining project. *BMC*
Plant Biol 2010, **10**:1471–2229.
- 529
530
531
532
41. Bhullar NK, Street K, Mackay M, Yahiaoui N, Keller B:
Unlocking wheat genetic resources for the molecular
identification of previously undescribed functional
alleles at the Pm3 resistance locus. *Proc Natl Acad Sci* 2009,
106:9519–9524.
- 533
534
535
536
537
42. Seeholzer S, Tsuchimatsu T, Jordan T, Bieri S, Pajonk S, Yang W,
Jahoor A, Shimizu KK, Keller B, Schulze-Lefert P: **Diversity at the**
Mla powdery mildew resistance locus from cultivated barley
reveals sites of positive selection. *Mol Plant–Microbe Interact*
2010, **23**:497–509.
- 538
539
540
541
43. Andersen EJ, Ali S, Reese RN, Yen Y, Neupane S: **Diversity and**
evolution of disease resistance genes in barley (*Hordeum*
vulgare L.). *Lib Acad* 2016 <http://dx.doi.org/10.4137/EBO.S38085>.TYPE.
- 542
543
544
545